

## **Transportation Infrastructure and Access on National Forests and Grasslands** **A Literature Review**

### **Introduction**

The Forest Service transportation system is very large with 374,883 miles (603,316 km) of system roads and 143,346 miles (230,693 km) of system trails. The system extends broadly across every national forest and grasslands and through a variety of habitats, ecosystems and terrains. An impressive body of scientific literature exists addressing the various effects of roads on the physical, biological and cultural environment – so much so, in the last few decades a new field of “road ecology” has emerged. In recent years, the scientific literature has expanded to address the effects of roads on climate change adaptation and conversely the effects of climate change on roads, as well as the effects of restoring lands occupied by roads on the physical, biological and cultural environments.

The following literature review summarizes the most recent thinking related to the environmental impacts of forest roads and motorized routes and ways to address them. The literature review is divided into three sections that address the environmental effects of transportation infrastructure on forests, climate change and infrastructure, and creating sustainable forest transportation systems.

- I. [Impacts of Transportation Infrastructure and Access to the Ecological Integrity of Terrestrial and Aquatic Ecosystems and Watersheds](#)
- II. [Climate Change and Transportation Infrastructure Including the Value of Roadless Areas for Climate Change Adaptation](#)
- III. [Sustainable Transportation Management in National Forests as Part of Ecological Restoration](#)

### **I. Impacts of Transportation Infrastructure and Access to the Ecological Integrity of Terrestrial and Aquatic Ecosystems and Watersheds**

It is well understood that transportation infrastructure and access management impact aquatic and terrestrial environments at multiple scales, and, in general, the more roads and motorized routes the greater the impact. In fact, in the past 20 years or so, scientists having realized the magnitude and breadth of ecological issues related to roads; entire books have been written on the topic, e.g., Forman et al. (2003), and a new scientific field called “road ecology” has emerged. Road ecology research centers have been created including the Western Transportation Institute at Montana State University and the Road Ecology Center at the University of California - Davis.<sup>1</sup>


---

<sup>1</sup> See <http://www.westerntransportationinstitute.org/research/roadecology> and <http://roadecology.ucdavis.edu/>

Below, we provide a summary of the current understanding on the impacts of roads and access allowed by road networks to terrestrial and aquatic ecosystems, drawing heavily on Gucinski et al. (2000). Other notable recent peer-reviewed literature reviews on roads include Trombulak and Frissell (2000), Switalski et al. (2004), Coffin (2007), Fahrig and Rytwinski (2009), and Robinson et al. (2010). Recent reviews on the impact of motorized recreation include Joslin and Youmans (1999), Gaines et al. (2003), Davenport and Switalski (2006), Ouren et al. (2007), and Switalski and Jones (2012). These peer-reviewed summaries provide additional information to help managers develop more sustainable transportation systems

### Impact on geomorphology and hydrology

The construction or presence of forest roads can dramatically change the hydrology and geomorphology of a forest system leading to reductions in the quantity and quality of aquatic habitat. While there are several mechanisms that cause these impacts (Wemple et al. 2001, Figure 1), most fundamentally, compacted roadbeds reduce rainfall infiltration, intercepting and concentrating water, and providing a ready source of sediment for transport (Wemple et al. 1996, Wemple et al. 2001). In fact, roads contribute more sediment to streams than any other land management activity (Gucinski et al. 2000). Surface erosion rates from roads are typically at least an order of magnitude greater than rates from harvested areas, and three orders of magnitude greater than erosion rates from undisturbed forest soils (Endicott 2008).



**Figure 1:** Typology of erosional and depositional features produced by mass-wasting and fluvial processes associate with forest roads (reprinted from Wemple et al. 2001)

Erosion of sediment from roads occurs both chronically and catastrophically. Every time it rains, sediment from the road surface and from cut- and fill-slopes is picked up by rainwater that flows into and on roads (fluvial erosion). The sediment that is entrained in surface flows are often concentrated into road ditches and culverts and directed into streams. The degree of fluvial erosion varies by geology and geography, and increases with increased motorized use (Robichaud et al. 2010). Closed roads produce less sediment, and Foltz et al. (2009) found a significant increase in erosion when closed roads were opened and driven upon.

Roads also precipitate catastrophic failures of road beds and fills (mass wasting) during large storm events leading to massive slugs of sediment moving into waterways (Endicott 2008; Gucinski et al. 2000). This typically occurs when culverts are undersized and cannot handle the volume of water, or they simply become plugged with debris. The saturated roadbed can fail entirely and result in a landslide, or the blocked stream crossing can erode the entire fill down to the original stream channel.

The erosion of road- and trail-related sediment and its subsequent movement into stream systems affects the geomorphology of the drainage system in a number of ways. The magnitude of their effects varies by climate, geology, road age, construction / maintenance practices and storm history. It directly alters channel morphology by embedding larger gravels as well as filling pools. It can also have the opposite effect of increasing peak discharges and scouring channels, which can lead to disconnection of the channel and floodplain, and lowered base flows (Furniss et al. 1991; Joslin and Youmans 1999). The width/depth ratio of the stream changes which then can trigger changes in water temperature, sinuosity and other geomorphic factors important for aquatic species survival (Joslin and Youmans 1999; Trombulak and Frissell 2000).

Roads also can modify flowpaths in the larger drainage network. Roads intercept subsurface flow as well as concentrate surface flow, which results in new flowpaths that otherwise would not exist, and the extension of the drainage network into previously unchanneled portions of the hillslope (Gucinski et al. 2000; Joslin and Youmans 1999). Severe aggradation of sediment at stream structures or confluences can force streams to actually go subsurface or make them too shallow for fish passage (Endicott 2008; Furniss et al. 1991).

### **Impacts on aquatic habitat and fish**

Roads can have dramatic and lasting impacts on fish and aquatic habitat. Increased sedimentation in stream beds has been linked to decreased fry emergence, decreased juvenile densities, loss of winter carrying capacity, and increased predation of fishes, and reductions in macro-invertebrate populations that are a food source to many fish species (Rhodes et al. 1994, Joslin and Youmans 1999, Gucinski et al. 2000, Endicott 2008). On a landscape scale, these effects can add up to: changes in the frequency, timing and magnitude of disturbance to aquatic habitat and changes to aquatic habitat structures (e.g., pools, riffles, spawning gravels and in-channel debris), and conditions (food sources, refugi, and water temperature) (Gucinski et al. 2000).

Roads can also act as barriers to migration (Gucinski et al. 2000). Where roads cross streams, road engineers usually place culverts or bridges. Culverts in particular can and often interfere with sediment transport and channel processes such that the road/stream crossing becomes a barrier for fish and aquatic species movement up and down stream. For instance, a culvert may scour on the downstream side of the crossing, actually forming a waterfall up which fish cannot

move. Undersized culverts and bridges can infringe upon the channel or floodplain and trap sediment causing the stream to become too shallow and/or warm such that fish will not migrate past the structure. This is problematic for many aquatic species but especially for anadromous species that must migrate upstream to spawn. Well-known native aquatic species affected by roads include salmon such as coho (*Oncorhynchus kisutch*), chinook (*O. tshawytscha*), and chum (*O. keta*); steelhead (*O. mykiss*); and a variety of trout species including bull trout (*Salvelinus confluentus*) and cutthroat trout (*O. clarki*), as well as other native fishes and amphibians (Endicott 2008).

### Impacts on terrestrial habitat and wildlife

Roads and trails impact wildlife through a number of mechanisms including: direct mortality (poaching, hunting/trapping) changes in movement and habitat use patterns (disturbance/avoidance), as well as indirect impacts including alteration of the adjacent habitat and interference with predatory/prey relationships (Wisdom et al. 2000, Trombulak and Frissell 2000). Some of these impacts result from the road itself, and some result from the uses on and around the roads (access). Ultimately, roads have been found to reduce the abundance and distribution of several forest species (Fayrig and Ritwinski 2009, Benítez-López et al. 2010).

**Table 1:** Road- and recreation trail-associated factors for wide-ranging carnivores (Reprinted from Gaines et al. (2003)<sup>2</sup>)

Focal species	Road-associated factors	Motorized trail-associated factors	Nonmotorized trail-associated factors
Grizzly bear	Poaching	Poaching	Poaching
	Collisions	Negative human interactions	Negative human interactions
	Negative human interactions	Displacement or avoidance	Displacement or avoidance
	Displacement or avoidance		
Lynx	Down log reduction	Disturbance at a specific site	Disturbance at a specific site
	Trapping	Trapping	
	Collisions		
	Disturbance at a specific site		
Gray wolf	Trapping	Trapping	Trapping
	Poaching	Disturbance at a specific site	Disturbance at a specific site
	Collisions		
	Negative human interactions		
	Disturbance at a specific site		
Wolverine	Displacement or avoidance		
	Down log reduction	Trapping	Trapping
	Trapping	Disturbance at a specific site	Disturbance at a specific site
	Disturbance at a specific site		

<sup>2</sup> For a list of citations see Gaines et al. (2003)

Direct mortality and disturbance from road and trail use impacts many different types of species. For example, wide-ranging carnivores can be significantly impacted by a number of factors including trapping, poaching, collisions, negative human interactions, disturbance and displacement (Gaines et al. 2003, Table 1). Hunted game species such as elk (*Cervus canadensis*), become more vulnerable from access allowed by roads and motorized trails resulting in a reduction in effective habitat among other impacts (Rowland et al. 2005, Switalski and Jones 2012). Slow-moving migratory animals such as amphibians, and reptiles who use roads to regulate temperature are also vulnerable (Gucinski et al. 2000, Brehme et al. 2013).

Habitat alteration is a significant consequence of roads as well. At the landscape scale, roads fragment habitat blocks into smaller patches that may not be able to support successfully interior forest species. Smaller habitat patches also results in diminished genetic variability, increased inbreeding, and at times local extinctions (Gucinski et al. 2000; Trombulak and Frissell 2000). Roads also change the composition and structure of ecosystems along buffer zones, called edge-affected zones. The width of edge-affected zones varies by what metric is being discussed; however, researchers have documented road-avoidance zones a kilometer or more away from a road (Table 2). In heavily roaded landscapes, edge-affected acres can be a significant fraction of total acres. For example, in a landscape area where the road density is 3 mi/mi<sup>2</sup> (not an uncommon road density in national forests) and where the edge-affected zone is estimated to be 500 ft from the center of the road to each side, the edge-affected zone is 56% of the total acreage.

**Table 2:** A summary of some documented road-avoidance zones for various species (adapted from Robinson et al. 2010).

Species	Avoidance zone		Reference
	m (ft)	Type of disturbance	
Snakes	650 (2133)	Forestry roads	Bowles (1997)
Salamander	35 (115)	Narrow forestry road, light traffic	Semlitsch (2003)
Woodland birds	150 (492)	Unpaved roads	Ortega and Capen (2002)
Spotted owl	400 (1312)	Forestry roads, light traffic	Wasser et al. (1997)
Marten	<100 (<328)	Any forest opening	Hargis et al. (1999)
Elk	500–1000 (1640-3281)	Logging roads, light traffic	Edge and Marcum (1985)
	100–300 (328-984)	Mountain roads depending on traffic volume	Rost and Bailey (1979)
Grizzly bear	3000 (9840)	Fall	Mattson et al. (1996)
	500 (1640)	Spring and summer	
	883 (2897)	Heavily traveled trail	Kasworm and Manley (1990)
	274 (899)	Lightly traveled trail	
	1122 (3681)	Open road	Kasworm and Manley (1990)
	665 (2182)	Closed road	

Black bear	274 (899)	Spring, unpaved roads	Kasworm and Manley (1990)
	914 (2999)	Fall, unpaved roads	

Roads and trails also affect ecosystems and habitats because they are also a major vector of non-native plant and animal species. This can have significant ecological and economic impacts when the invading species are aggressive and can overwhelm or significantly alter native species and systems. In addition, roads can increase harassment, poaching and collisions with vehicles, all of which lead to stress or mortality (Wisdom et al. 2000).

Recent reviews have synthesized the impacts of roads on animal abundance and distribution. Fahrig and Rytwinski (2009) did a complete review of the empirical literature on effects of roads and traffic on animal abundance and distribution looking at 79 studies that addressed 131 species and 30 species groups. They found that the number of documented negative effects of roads on animal abundance outnumbered the number of positive effects by a factor of 5. Amphibians, reptiles, most birds tended to show negative effects. Small mammals generally showed either positive effects or no effect, mid-sized mammals showed either negative effects or no effect, and large mammals showed predominantly negative effects. Benítez-López et al. (2010) conducted a meta-analysis on the effects of roads and infrastructure proximity on mammal and bird populations. They found a significant pattern of avoidance and a reduction in bird and mammal populations in the vicinity of infrastructure.

### Road density<sup>3</sup> thresholds for fish and wildlife

It is well documented that beyond specific road density thresholds, certain species will be negatively affected, and some will be extirpated. Most studies that look into the relationship between road density and wildlife focus on the impacts to large endangered carnivores or hunted game species, although high road densities certainly affect other species – for instance, reptiles and amphibians. Gray wolves (*Canis lupus*) in the Great Lakes region and elk in Montana and Idaho have undergone the most long-term and in depth analysis. Forman and Hersperger (1996) found that in order to maintain a naturally functioning landscape with sustained populations of large mammals, road density must be below 0.6 km/km<sup>2</sup> (1.0 mi/mi<sup>2</sup>). Several studies have since substantiated their claim (Robinson et al. 2010, Table 3).

A number of studies at broad scales have also shown that higher road densities generally lead to greater impacts to aquatic habitats and fish density (Table 3). Carnefix and Frissell (2009) provide a concise review of studies that correlate cold water fish abundance and road density, and from the cited evidence concluded that “1) no truly “safe” threshold road density exists, but rather negative impacts begin to accrue and be expressed with incursion of the very first road segment; and 2) highly significant impacts (e.g., threat of extirpation of sensitive species) are already apparent at road densities on the order of 0.6 km/km<sup>2</sup> (1.0 mi/mi<sup>2</sup>) or less” (p. 1).

<sup>3</sup> We intend the term “road density” to refer to the density all roads within national forests, including system roads, closed roads, non-system roads administered by other jurisdictions (private, county, state), temporary roads and motorized trails. Please see Attachment 2 for the relevant existing scientific information supporting this approach.

**Table 3:** A summary of some road-density thresholds and correlations for terrestrial and aquatic species and ecosystems (reprinted from Robinson et al. 2010).

Species (Location)	Road density (mean, guideline, threshold, correlation)	Reference
Wolf (Minnesota)	0.36 km/km <sup>2</sup> (mean road density in primary range); 0.54 km/km <sup>2</sup> (mean road density in peripheral range)	Mech et al. (1988)
Wolf	>0.6 km/km <sup>2</sup> (absent at this density)	Jalkotzy et al. (1997)
Wolf (Northern Great Lakes region)	>0.45 km/km <sup>2</sup> (few packs exist above this threshold); >1.0 km/km <sup>2</sup> (no pack exist above this threshold)	Mladenoff et al. (1995)
Wolf (Wisconsin)	0.63 km/km <sup>2</sup> (increasing due to greater human tolerance)	Wydeven et al. (2001)
Wolf, mountain lion (Minnesota, Wisconsin, Michigan)	0.6 km/km <sup>2</sup> (apparent threshold value for a naturally functioning landscape containing sustained populations)	Thiel (1985); van Dyke et al. (1986); Jensen et al. (1986); Mech et al. (1988); Mech (1989)
Elk (Idaho)	1.9 km/km <sup>2</sup> (density standard for habitat effectiveness)	Woodley 2000 cited in Beazley et al. 2004
Elk (Northern US)	1.24 km/km <sup>2</sup> (habitat effectiveness decline by at least 50%)	Lyon (1983)
Elk, bear, wolverine, lynx, and others	0.63 km/km <sup>2</sup> (reduced habitat security and increased mortality)	Wisdom et al. (2000)
Moose (Ontario)	0.2-0.4 km/km <sup>2</sup> (threshold for pronounced response)	Beyer et al. (2013)
Grizzly bear (Montana)	>0.6 km/km <sup>2</sup>	Mace et al. (1996); Mattson et al. (1996)
Black bear (North Carolina)	>1.25 km/km <sup>2</sup> (open roads); >0.5 km/km <sup>2</sup> (logging roads); (interference with use of habitat)	Brody and Pelton (1989)
Black bear	0.25 km/km <sup>2</sup> (road density should not exceed)	Jalkotzy et al. (1997)
Bobcat (Wisconsin)	1.5 km/km <sup>2</sup> (density of all road types in home range)	Jalkotzy et al. (1997)
Large mammals	>0.6 km/km <sup>2</sup> (apparent threshold value for a naturally functioning landscape containing sustained populations)	Forman and Hersperger (1996)
Bull trout (Montana)	Inverse relationship of population and road density	Rieman et al. (1997); Baxter et al. (1999)
Fish populations (Medicine Bow National Forest)	(1) Positive correlation of numbers of culverts and stream crossings and amount of fine sediment in stream channels (2) Negative correlation of fish density and numbers of culverts	Eaglin and Hubert (1993) cited in Gucinski et al. (2001)
Macroinvertebrates	Species richness negatively correlated with an index of	McGurk and Fong (1995)

	road density	
Non-anadromous salmonids (Upper Columbia River basin)	(1) Negative correlation likelihood of spawning and rearing and road density (2) Negative correlation of fish density and road density	Lee et al. (1997)

Where both stream and road densities are high, the incidence of connections between roads and streams can also be expected to be high, resulting in more common and pronounced effects of roads on streams (Gucinski et al. 2000). For example, a study on the Medicine Bow National Forest (WY) found as the number of culverts and stream crossings increased, so did the amount of sediment in stream channels (Eaglin and Hubert 1993). They also found a negative correlation with fish density and the number of culverts. Invertebrate communities can also be impacted. McGurk and Fong (1995) report a negative correlation between an index of road density with macroinvertebrate diversity.

The U.S. Fish and Wildlife Service’s Final Rule listing bull trout as threatened (USDI Fish and Wildlife Service 1999) addressed road density, stating:

“... assessment of the interior Columbia Basin ecosystem revealed that increasing road densities were associated with declines in four non-anadromous salmonid species (bull trout, Yellowstone cutthroat trout, westslope cutthroat trout, and redband trout) within the Columbia River Basin, likely through a variety of factors associated with roads (Quigley & Arbelbide 1997). Bull trout were less likely to use highly roaded basins for spawning and rearing, and if present, were likely to be at lower population levels (Quigley and Arbelbide 1997). Quigley et al. (1996) demonstrated that when average road densities were between 0.4 to 1.1 km/km<sup>2</sup> (0.7 and 1.7 mi/mi<sup>2</sup>) on USFS lands, the proportion of subwatersheds supporting “strong” populations of key salmonids dropped substantially. Higher road densities were associated with further declines” (USDI Fish and Wildlife Service 1999, p. 58922).

Anderson et al. (2012) also showed that watershed conditions tend to be best in areas protected from road construction and development. Using the US Forest Service’s Watershed Condition Framework assessment data, they showed that National Forest lands that are protected under the Wilderness Act, which provides the strongest safeguards, tend to have the healthiest watersheds. Watersheds in Inventoried Roadless Areas – which are protected from road building and logging by the Roadless Area Conservation Rule – tend to be less healthy than watersheds in designated Wilderness, but they are considerably healthier than watersheds in the managed landscape.

### Impacts on other resources

Roads and motorized trails also play a role in affecting wildfire occurrence. Research shows that human-ignited wildfires, which account for more than 90% of fires on national lands, is almost five times more likely in areas with roads (USDA Forest Service 1996a; USDA Forest Service 1998). Furthermore, Baxter (2002) found that off-road vehicles (ORVs) can be a significant source of fire ignitions on forestlands. Roads can affect where and how forests burn and, by extension, the vegetative condition of the forest. See Attachment 1 for more information documenting the relationship between roads and wildfire occurrence.

Finally, access allowed by roads and trails can increase of ORV and motorized use in remote areas threatening archaeological and historic sites. Increased visitation has resulted in



intentional and unintentional damage to many cultural sites (USDI Bureau of Land Management 2000, Schiffman 2005).

## **II. Climate Change and Transportation Infrastructure including the value of roadless areas for climate change adaptation**

As climate change impacts grow more profound, forest managers must consider the impacts on the transportation system as well as from the transportation system. In terms of the former, changes in precipitation and hydrologic patterns will strain infrastructure at times to the breaking point resulting in damage to streams, fish habitat, and water quality as well as threats to public safety. In terms of the latter, the fragmenting effect of roads on habitat will impede the movement of species which is a fundamental element of adaptation. Through planning, forest managers can proactively address threats to infrastructure, and can actually enhance forest resilience by removing unneeded roads to create larger patches of connected habitat.

### **Impact of climate change and roads on transportation infrastructure**

It is expected that climate change will be responsible for more extreme weather events, leading to increasing flood severity, more frequent landslides, changing hydrographs (peak, annual mean flows, etc.), and changes in erosion and sedimentation rates and delivery processes. Roads and trails in national forests, if designed by an engineering standard at all, were designed for storms and water flows typical of past decades, and hence may not be designed for the storms in future decades. Hence, climate driven changes may cause transportation infrastructure to malfunction or fail (ASHTO 2012, USDA Forest Service 2010). The likelihood is higher for facilities in high-risk settings—such as rain-on-snow zones, coastal areas, and landscapes with unstable geology (USDA Forest Service 2010).

Forests fragmented by roads will likely demonstrate less resistance and resilience to stressors, like those associated with climate change (Noss 2001). First, the more a forest is fragmented (and therefore the higher the edge/interior ratio), the more the forest loses its inertia characteristic, and becoming less resilient and resistant to climate change. Second, the more a forest is fragmented characterized by isolated patches, the more likely the fragmentation will interfere with the ability of species to track shifting climatic conditions over time and space. Noss (2001) predicts that weedy species with effective dispersal mechanisms might benefit from fragmentation at the expense of native species.

### **Modifying infrastructure to increase resilience**

To prevent or reduce road failures, culvert blow-outs, and other associated hazards, forest managers will need to take a series of actions. These include replacing undersized culverts with larger ones, prioritizing maintenance and upgrades (e.g., installing drivable dips and more outflow structures), and obliterating roads that are no longer needed and pose erosion hazards (USDA Forest Service 2010, USDA Forest Service 2012a, USDA Forest Service 2011, Table 4).

Olympic National Forest has developed a number of documents oriented at oriented at protecting watershed health and species in the face of climate change, including a 2003 travel management strategy and a report entitled *Adapting to Climate Change in Olympic National Park and National Forest*. In the travel management strategy, Olympic National Forest recommended that 1/3<sup>rd</sup> of its road system be decommissioned and obliterated (USDA Forest Service 2011a). In addition, the plan called for addressing fish migration barriers in a prioritized and strategic way – most of these are associated with roads. The report calls for road decommissioning, relocation of roads away from streams, enlarging culverts as well as replacing culverts with fish-friendly crossings (USDA Forest Service 2011a, Table 4).

**Table 4:** Current and expected sensitivities of fish to climate change on the Olympic Peninsula, associated adaptation strategies and action for fisheries and fish habitat management and relevant to transportation management at Olympic National Forest and Olympic National Park (excerpt reprinted from USDA Forest Service 2011a).

Current and expected sensitivities	Adaptation strategies and actions
Changes in habitat quantity and quality	<ul style="list-style-type: none"> <li>• Implement habitat restoration projects that focus on re-creating watershed processes and functions and that create diverse, resilient habitat.</li> </ul>
Increase in culvert failures, fill-slope failures, stream adjacent road failures, and encroachment from stream-adjacent road segments	<ul style="list-style-type: none"> <li>• Decommission unneeded roads.</li> <li>• Remove sidecast, improve drainage, and increase culvert sizing on remaining roads.</li> <li>• Relocate stream-adjacent roads.</li> </ul>
Greater difficulty disconnecting roads from stream channels	<ul style="list-style-type: none"> <li>• Design more resilient stream crossing structures.</li> </ul>
Major changes in quantity and timing of streamflow in transitional watersheds	<ul style="list-style-type: none"> <li>• Make road and culvert designs more conservative in transitional watersheds to accommodate expected changes.</li> </ul>
Decrease in area of headwater streams	<ul style="list-style-type: none"> <li>• Continue to correct culvert fish passage barriers.</li> <li>• Consider re-prioritizing culvert fish barrier correction projects.</li> </ul>
Decrease in habitat quantity and connectivity for species that use headwater streams	<ul style="list-style-type: none"> <li>• Restore habitat in degraded headwater streams that are expected to retain adequate summer streamflow (ONF).</li> </ul>

In December 2012, the USDA Forest Service published a report entitled “Assessing the Vulnerability of Watersheds to Climate Change.” This document reinforces the concept expressed by Olympic National Forest that forest managers need to be proactive in reducing erosion potential from roads:

“Road improvements were identified as a key action to improve condition and resilience of watersheds on all the pilot Forests. In addition to treatments that reduce erosion, road improvements can reduce the delivery of runoff from road segments to channels, prevent diversion of flow during large events, and restore aquatic habitat connectivity by providing for passage of aquatic organisms. As stated previously, watershed sensitivity is determined by both inherent and management-related factors. Managers have no control over the inherent factors,

so to improve resilience, efforts must be directed at anthropogenic influences such as instream flows, roads, rangeland, and vegetation management....

[Watershed Vulnerability Analysis] results can also help guide implementation of travel management planning by informing priority setting for decommissioning roads and road reconstruction/maintenance. As with the Ouachita NF example, disconnecting roads from the stream network is a key objective of such work. Similarly, WVA analysis could also help prioritize aquatic organism passage projects at road-stream crossings to allow migration by aquatic residents to suitable habitat as streamflow and temperatures change" (USDA Forest Service 2012a, p. 22-23).

### **Reducing fragmentation to enhance aquatic and terrestrial species adaptation**

Decommissioning and upgrading roads and thus reducing the amount of fine sediment deposited on salmonid nests can increase the likelihood of egg survival and spawning success (McCaffery et al. 2007). In addition, this would reconnect stream channels and remove barriers such as culverts. Decommissioning roads in riparian areas may provide further benefits to salmon and other aquatic organisms by permitting reestablishment of streamside vegetation, which provides shade and maintains a cooler, more moderated microclimate over the stream (Battin et al. 2007).

One of the most well documented impacts of climate change on wildlife is a shift in the ranges of species (Parmesan 2006). As animals migrate, landscape connectivity will be increasingly important (Holman et al. 2005). Decommissioning roads in key wildlife corridors will improve connectivity and be an important mitigation measure to increase resiliency of wildlife to climate change. For wildlife, road decommissioning can reduce the many stressors associated with roads. Road decommissioning restores habitat by providing security and food such as grasses and fruiting shrubs for wildlife (Switalski and Nelson 2011).

Forests fragmented by roads and motorized trail networks will likely demonstrate less resistance and resilience to stressors, such as weeds. As a forest is fragmented and there is more edge habitat, Noss (2001) predicts that weedy species with effective dispersal mechanisms will increasingly benefit at the expense of native species. However, decommissioned roads when seeded with native species can reduce the spread of invasive species (Grant et al. 2011), and help restore fragmented forestlands. Off-road vehicles with large knobby tires and large undercarriages are also a key vector for weed spread (e.g., Rooney 2006). Strategically closing and decommissioning motorized routes, especially in roadless areas, will reduce the spread of weeds on forestlands (Gelbard and Harrison 2003).

### **Transportation infrastructure and carbon sequestration**

The topic of the relationship of road restoration and carbon has only recently been explored. There is the potential for large amounts of carbon (C) to be sequestered by reclaiming roads. When roads are decompacted during reclamation, vegetation and soils can develop more rapidly and sequester large amounts of carbon. A recent study estimated total soil C storage increased 6 fold to 6.5 x 107g C/km (to 25 cm depth) in the northwestern US compared to untreated abandoned roads (Lloyd et al. 2013). Another recent study concluded that reclaiming 425 km of logging roads over the last 30 years in Redwood National Park in Northern California resulted in net carbon savings of 49,000 Mg carbon to date (Madej et al. 2013, Table 5).

Kerekvliet et al. (2008) published a Wilderness Society briefing memo on the impact to carbon sequestration from road decommissioning. Using Forest Service estimates of the fraction of road miles that are unneeded, the authors calculated that restoring 126,000 miles of roads to a natural state would be equivalent to revegetating an area larger than Rhode Island. In addition, they calculate that the net economic benefit of road treatments are always positive and range from US\$0.925-1.444 billion.

**Table 5.** Carbon budget implications in road decommissioning projects (reprinted from Madej et al. 2013).

Road Decommissioning Activities and Processes	Carbon Cost	Carbon Savings
Transportation of staff to restoration sites (fuel emissions)	X	
Use of heavy equipment in excavations (fuel emissions)	X	
Cutting trees along road alignment during hillslope recontouring	X	
Excavation of road fill from stream crossings		X
Removal of road fill from unstable locations		X
Reduces risk of mass movement		X
Post-restoration channel erosion at excavation sites	X	
Natural revegetation following road decompaction		X
Replanting trees		X
Soil development following decompaction		X

**Benefits of roadless areas and roadless area networks to climate change adaptation**

Undeveloped natural lands provide numerous ecological benefits. They contribute to biodiversity, enhance ecosystem representation, and facilitate connectivity (Loucks et al. 2003; Crist and Wilmer 2002, Wilcove 1990, The Wilderness Society 2004, Strittholt and Dellasala 2001, DeVelice and Martin 2001), and provide high quality or undisturbed water, soil and air (Anderson et al. 2012, Dellasalla et al. 2011). They also can serve as ecological baselines to help us better understand our impacts to other landscapes, and contribute to landscape resilience to climate change.

Forest Service roadless lands, in particular, are heralded for the conservation values they provide. These are described at length in the preamble of the Roadless Area Conservation Rule (RACR)<sup>4</sup> as well as in the Final Environmental Impact Statement (FEIS) for the RACR<sup>5</sup>, and include: high quality or undisturbed soil, water, and air; sources of public drinking water; diversity of plant and animal communities; habitat for threatened, endangered, proposed, candidate, and sensitive species and for those species dependent on large, undisturbed areas of land; primitive, semi-primitive non- motorized, and semi-primitive motorized classes of

<sup>4</sup> Federal Register .Vol. 66, No. 9. January 12, 2001. Pages 3245-3247.

<sup>5</sup> Final Environmental Impact Statement, Vol. 1, 3-3 to 3-7

dispersed recreation; reference landscapes; natural appearing landscapes with high scenic quality; traditional cultural properties and sacred sites; and other locally identified unique characteristics (e.g., include uncommon geological formations, unique wetland complexes, exceptional hunting and fishing opportunities).

The Forest Service, National Park Service, and US Fish and Wildlife Service recognize that protecting and connecting roadless or lightly roaded areas is an important action agencies can take to enhance climate change adaptation. For example, the Forest Service National Roadmap for Responding to Climate Change (USDA Forest Service 2011b) establishes that increasing connectivity and reducing fragmentation are short and long term actions the Forest Service should take to facilitate adaptation to climate change.<sup>6</sup> The National Park Service also identifies connectivity as a key factor for climate change adaptation along with establishing “blocks of natural landscape large enough to be resilient to large-scale disturbances and long-term changes” and other factors. The agency states that: “The success of adaptation strategies will be enhanced by taking a broad approach that identifies connections and barriers across the landscape. Networks of protected areas within a larger mixed landscape can provide the highest level of resilience to climate change.”<sup>7</sup> Similarly, the National Fish, Wildlife and Plants Climate Adaptation Partnership’s Adaptation Strategy (2012) calls for creating an ecologically-connected network of conservation areas.<sup>8</sup>

Crist and Wilmer (2002) looked at the ecological value of roadless lands in the Northern Rockies and found that protection of national forest roadless areas, when added to existing federal conservation lands in the study area, would 1) increase the representation of virtually all land cover types on conservation lands at both the regional and ecosystem scales, some by more

---

<sup>6</sup> Forest Service, 2011. *National Roadmap for Responding to Climate Change*. US Department of Agriculture. FS-957b. Page 26.

<sup>7</sup> National Park Service. *Climate Change Response Program Brief*. <http://www.nature.nps.gov/climatechange/adaptationplanning.cfm>. Also see: National Park Service, 2010. *Climate Change Response Strategy*. [http://www.nature.nps.gov/climatechange/docs/NPS\\_CCRS.pdf](http://www.nature.nps.gov/climatechange/docs/NPS_CCRS.pdf). Objective 6.3 is to “Collaborate to develop cross-jurisdictional conservation plans to protect and restore connectivity and other landscape-scale components of resilience.”

<sup>8</sup> See <http://www.wildlifeadaptationstrategy.gov/pdf/NFWPCAS-Chapter-3.pdf>. Pages 55- 59. The first goal and related strategies are:

Goal 1: Conserve habitat to support healthy fish, wildlife, and plant populations and ecosystem functions in a changing climate.

Strategy 1.1: identify areas for an ecologically-connected network of terrestrial, freshwater, coastal, and marine conservation areas that are likely to be resilient to climate change and to support a broad range of fish, wildlife, and plants under changed conditions.

Strategy 1.2: Secure appropriate conservation status on areas identified in Strategy 1.1 to complete an ecologically-connected network of public and private conservation areas that will be resilient to climate change and support a broad range of species under changed conditions.

Strategy 1.4: Conserve, restore, and as appropriate and practicable, establish new ecological connections among conservation areas to facilitate fish, wildlife, and plant migration, range shifts, and other transitions caused by climate change.

than 100%; 2) help protect rare, species-rich, and often-declining vegetation communities; and 3) connect conservation units to create bigger and more cohesive habitat “patches.”

Roadless lands also are responsible for higher quality water and watersheds. Anderson et al. (2012) assessed the relationship of watershed condition and land management status and found a strong spatial association between watershed health and protective designations. Dellasalla et al. (2011) found that undeveloped and roadless watersheds are important for supplying downstream users with high-quality drinking water, and developing these watersheds comes at significant costs associated with declining water quality and availability. The authors recommend a light-touch ecological footprint to sustain the many values that derive from roadless areas including healthy watersheds.

### **III. Sustainable Transportation Management in National Forests as Part of Ecological Restoration**

At 375,000 miles strong, the Forest Service road system is one of the largest in the world – it is eight times the size of the National Highway System. It is also indisputably unsustainable – that is, roads are not designed, located, or maintained according to best management practices, and environmental impacts are not minimized. It is largely recognized that forest roads, especially unpaved ones, are a primary source of sediment pollution to surface waters (Endicott 2008, Gucinski et al. 2000), and that the system has about 1/3<sup>rd</sup> more miles than it needs (USDA Forest Service 2001). In addition, the majority of the roads were constructed decades ago when road design and management techniques did not meet current standards (Gucinski et al. 2000, Endicott 2008), making them more vulnerable to erosion and decay than if they had been designed today. Road densities in national forests often exceed accepted thresholds for wildlife.

Only a small portion of the road system is regularly used. All but 18% of the road system is inaccessible to passenger vehicles. Fifty-five percent of the roads are accessible only by high clearance vehicles and 27% are closed. The 18% that is accessible to cars is used for about 80% of the trips made within National Forests.<sup>9</sup> Most of the road maintenance funding is directed to the passenger car roads, while the remaining roads suffer from neglect. As a result, the Forest Service currently has a \$3.7 billion road maintenance backlog that grows every year. In other words, only about 1/5th of the roads in the national forest system are used most of the time, and the fraction that is used often is the best designed and maintained because they are higher level access roads. The remaining roads sit generally unneeded and under-maintained – arguably a growing ecological and fiscal liability.

Current Forest Service management direction is to identify and implement a sustainable transportation system.<sup>10</sup> The challenge for forest managers is figuring out what is a sustainable road system and how to achieve it – a challenge that is exacerbated by climate change. It is reasonable to define a sustainable transportation system as one where all the routes are constructed, located, and maintained with best management practices, and social and environmental impacts are minimized. This, of course, is easier said than done, since the reality

---

<sup>9</sup> USDA Forest Service. Road Management Website Q&As. Available online at [http://www.fs.fed.us/eng/road\\_mgt/qanda.shtml](http://www.fs.fed.us/eng/road_mgt/qanda.shtml).

<sup>10</sup> See Forest Service directive memo dated March 29, 2012 entitled “Travel Management, Implementation of 36 CFR, Part 202, Subpart A (36 CFR 212.5(b))”

is that even the best roads and trail networks can be problematic simply because they exist and usher in land uses that without the access would not occur (Trombulak and Frissell 2000, Carnefix and Frissell 2009, USDA Forest Service 1996b), and when they are not maintained to the designed level they result in environmental problems (Endicott 2008; Gucinski et al. 2000). Moreover, what was sustainable may no longer be sustainable under climate change since roads designed to meet older climate criteria may no longer hold up under new climate scenarios (USDA Forest Service 2010, USDA Forest Service 2011b, USDA Forest Service 2012a, AASHTO 2012).

### **Forest Service efforts to move toward a more sustainable transportation system**

The Forest Service has made efforts to make its transportation system more sustainable, but still has considerable work to do. In 2001, the Forest Service tried to address the issue by promulgating the Roads Rule<sup>11</sup> with the purpose of working toward a sustainable road system (USDA 2001). The Rule directed every national forest to identify a minimum necessary road system and identify unneeded roads for decommissioning. To do this, the Forest Service developed the Roads Analysis Process (RAP), and published Gucinski et al. (2000) to provide the scientific foundation to complement the RAP. In describing the RAP, Gucinski et al. (2000) writes:

“Roads Analysis is intended to be an integrated, ecological, social, and economic approach to transportation planning. It uses a multiscale approach to ensure that the identified issues are examined in context. Roads Analysis is to be based on science. Analysts are expected to locate, correctly interpret, and use relevant existing scientific literature in the analysis, disclose any assumptions made during the analysis, and reveal the limitations of the information on which the analysis is based. The analysis methods and the report are to be subjected to critical technical review” (p. 10).

Most national forests have completed RAPs, although most only looked at passenger vehicle roads which account for less than 20% of the system’s miles. The Forest Service Washington Office in 2010 directed that forests complete a Travel Analysis Process (TAP) by the end of fiscal year 2015, which must address all roads and create a map and list of roads identifying which are likely needed and which are not. Completed TAPs will provide a blueprint for future road decommissioning and management, they will not constitute compliance with the Roads Rule, which clearly requires the identification of the minimum roads system and roads for decommissioning. Almost all forests have yet to comply with subpart A.

The Forest Service in 2005 then tried to address the off-road portion of this issue by promulgating subpart B of the Travel Management Rule,<sup>12</sup> with the purpose of curbing the most serious impacts associated with off-road vehicle use. Without a doubt, securing summer-time travel management plans was an important step to curbing the worst damage. However, much work remains to be done to approach sustainability, especially since many national forests used the travel management planning process to simply freeze the footprint of motorized routes, and did not try to re-design the system to make it more ecologically or socially sustainable. Adams and McCool (2009) considered this question of how to achieve sustainable motorized recreation and concluded that:

---

<sup>11</sup> 36 CFR 215 subpart A

<sup>12</sup> 36 CFR 212 subpart B

As the agencies move to revise [off-road vehicle] allocations, they need to clearly define how they intend to locate routes so as to minimize impacts to natural resources and other recreationists in accordance with Executive Order 11644....<sup>13</sup>

...As they proceed with designation, the FS and BLM need to acknowledge that current allocations are the product of agency failure to act, not design. Ideally, ORV routes would be allocated as if the map were currently empty of ORV routes. Reliance on the current baseline will encourage inefficient allocations that likely disproportionately impact natural resources and non-motorized recreationists. While acknowledging existing use, the agencies need to do their best to imagine the best possible arrangement of ORV routes, rather than simply tinkering around the edges of the current allocations.<sup>14</sup>

The Forest Service only now is contemplating addressing the winter portion of the issue, forced by a lawsuit challenging the Forest Service's inadequate management of snowmobiles. The agency is expected to issue a third rule in the fall of 2014 that will trigger winter travel management planning.

### **Strategies for identifying a minimum road system and prioritizing restoration**

Transportation Management plays an integral role in the restoration of Forestlands. Reclaiming and obliterating roads is key to developing a sustainable transportation system. Numerous authors have suggested removing roads 1) to restore water quality and aquatic habitats (Gucinski et al. 2000), and 2) to improve habitat security and restore terrestrial habitat (e.g., USDI USFWS 1993, Hebblewhite et al. 2009).

Creating a minimum road system through road removal will increase connectivity and decrease fragmentation across the entire forest system. However, at a landscape scale, certain roads and road segments pose greater risks to terrestrial and aquatic integrity than others. Hence, restoration strategies must focus on identifying and removing/mitigating the higher risk roads. Additionally, areas with the highest ecological values, such as being adjacent to a roadless area, may also be prioritized for restoration efforts. Several methods have been developed to help prioritize road reclamation efforts including GIS-based tools and best management practices (BMPs). It is our hope that even with limited resources, restoration efforts can be prioritized and a more sustainable transportation system created.

### **GIS-based tools**

---

<sup>13</sup> Recent court decisions have made it clear that the minimization requirements in the Executive Orders are not discretionary and that the Executive Orders are enforceable. See


- *Idaho Conservation League v. Guzman*, 766 F. Supp. 2d 1056 (D. Idaho 2011) (Salmon-Challis National Forest TMP).
- *The Wilderness Society v. U.S. Forest Service*, CV 08-363 (D. Idaho 2012) (Sawtooth-Minidoka district National Forest TMP).
- *Central Sierra Environmental Resource Center v. US Forest Service*, CV 10-2172 (E.D. CA 2012) (Stanislaus National Forest TMP).

<sup>14</sup> Page 105.



Girvetz and Shilling (2003) developed a novel and inexpensive way to analyze environmental impacts from road systems using the Ecosystem Management Decision Support program (EMDS). EMDS was originally developed by the United States Forest Service, as a GIS-based decision support tool to conduct ecological analysis and planning (Reynolds 1999). Working in conjunction with Tahoe National Forest managers, Girvetz and Shilling (2003) used spatial data on a number of aquatic and terrestrial variables and modeled the impact of the forest's road network. The network analysis showed that out of 8233 km of road analyzed, only 3483 km (42%) was needed to ensure current and future access to key points. They found that the modified network had improved patch characteristics, such as significantly fewer "cherry stem" roads intruding into patches, and larger roadlessness.

Shilling et al. (2012) later developed a recreational route optimization model using a similar methodology and with the goal of identifying a sustainable motorized transportation system for the Tahoe National Forest (Figure 2). Again using a variety of environmental factors, the model identified routes with high recreational benefits, lower conflict, lower maintenance and management requirements, and lower potential for environmental impact operating under the presumption that such routes would be more sustainable and preferable in the long term. The authors combined the impact and benefit analyses into a recreation system analysis "that was effectively a cost-benefit accounting, consistent with requirements of both the federal Travel Management Rule (TMR) and the National Environmental Policy Act" (p. 392).



**Figure 2:** A knowledge base of contributions of various environmental conditions to the concept “environmental impact” [of motorized trails]. Rectangles indicate concepts, circles indicate Boolean logic operators, and rounded rectangles indicate sources of environmental data. (Reprinted from Shilling et al. 2012)

The Wilderness Society in 2012 also developed a GIS decision support tool called “RoadRight” that identifies high risk road segments to a variety of forest resources including water, wildlife, and roadlessness (The Wilderness Society 2012, The Wilderness Society 2013). The GIS system is designed to provide information that will help forest planners identify and minimize road related environmental risks. See the summary of and user guide for RoadRight that provides more information including where to access the open source software.<sup>15</sup>

<sup>15</sup> The Wilderness Society, 2012. Rightsizing the National Forest Road System: A Decision Support Tool. Available at <http://www.landscapecollaborative.org/download/attachments/12747016/Road+decommissioning+model+overview+2012-02-29.pdf?version=1&modificationDate=1331595972330>.

The Wilderness Society, 2013.  
RoadRight: A Spatial Decision Support System to Prioritize Decommissioning and Repairing Roads in

### **Best management practices (BMPs)**

BMPs have also been developed to help create more sustainable transportation systems and identify restoration opportunities. BMPs provide science-based criteria and standards that land managers follow in making and implementing decisions about human uses and projects that affect natural resources. Several states have developed BMPs for road construction, maintenance and decommissioning practices (e.g., Logan 2001, Merrill and Cassaday 2003, USDA Forest Service 2012b).

Recently, BMPs have been developed for addressing motorized recreation. Switalski and Jones (2012) published, "*Off-Road Vehicle Best Management Practices for Forestlands: A Review of Scientific Literature and Guidance for Managers.*" This document reviews the current literature on the environmental and social impacts of off-road vehicles (ORVs), and establishes a set of Best Management Practices (BMPs) for the planning and management of ORV routes on forestlands. The BMPs were designed to be used by land managers on all forestlands, and is consistent with current forest management policy and regulations. They give guidance to transportation planners on where how to place ORV routes in areas where they will reduce use conflicts and cause as little harm to the environment as possible. These BMPs also help guide managers on how to best remove and restore routes that are redundant or where there is an unacceptable environmental or social cost.

### **References**

- AASHTO. 2012. Adapting Infrastructure to Extreme Weather Events: Best Practices and Key Challenges. Background Paper. AASHTO Workshop. Traverse City, Michigan, May 20, 2012. Available at: [http://climatechange.transportation.org/pdf/adapt\\_background5-20-12.pdf](http://climatechange.transportation.org/pdf/adapt_background5-20-12.pdf).
- Adams, J.C., and S.F. McCool. 2009. Finite recreation opportunities: The Forest Service, the Bureau of Land Management, and off-road vehicle management. *Natural Areas Journal* 49: 45–116.
- Anderson, H.M., C. Gaolach, J. Thomson, and G. Aplet. 2012. Watershed Health in Wilderness, Roadless, and Roded Areas of the National Forest System. *Wilderness Society Report*. 11 p.

---

National Forests User Guide. RoadRight version: 2.2, User Guide version: February, 2013. Available at <http://www.landscapecollaborative.org/download/attachments/18415665/RoadRight%20User%20Guide%20v22.pdf?api=v2>

- Battin J., M.W. Wiley, M.H. Ruckelshaus, R.N. Palmer, E. Korb, K.K. Bartz, and H. Imaki. 2007. Projected impacts of climate change on salmon habitat restoration. *Proceedings of the National Academy of Sciences of the United States of America* 104: 6720–6725.
- Baxter, C.V., C.A. Frissell, and F.R. Hauer. 1999. Geomorphology, logging roads, and the distribution of bull trout spawning in a forested river basin: implications for management and conservation. *Transactions of the American Fisheries Society* 128: 854–867.
- Baxter, G. 2002. All terrain vehicles as a cause of fire ignition in Alberta forests. *Advantage* (Publication of the Forest Engineering Research Institute of Canada) 3(44): 1-7.
- Beazley, K., T. Snaith, F. MacKinnon, and D. Colville. 2004. Road density and the potential impacts on wildlife species such as American moose in mainland Nova Scotia. *Proceedings of the Nova Scotia Institute of Science* 42: 339-357.
- Benítez-López, A., R. Alkemade, and P.A. Verweij. 2010. The impacts of roads and other infrastructure on mammal and bird populations: a meta-analysis. *Biological Conservation* 143: 1307-1316.
- Beyer, H.L., R. Ung, D.L. Murray, and M.J. Fortin. 2013. Functional responses, seasonal variation and thresholds in behavioural responses of moose to road density. *Journal of Applied Ecology* 50: 286–294.
- Brehme, C.S., and J.A. Tracey, L.R. McClenaghan, and R.N. Fisher. 2013. Permeability of roads to movement of scrubland lizards and small mammals. *Conservation Biology* 27(4): 710–720.
- Bowles, A.E. 1997. Responses of wildlife to noise. In *Wildlife and recreationists: coexistence through management and research*. Edited by R.L. Knight and K.J. Gutzwiller. Island Press, Washington, DC. p. 109–156.
- Brody, A.J., and M.R. Pelton. 1989. Effects of roads on black bear movements in western North Carolina. *Wildlife Society Bulletin* 17: 5-10.
- Carnefix, G., and C. A. Frissell. 2009. Aquatic and Other Environmental Impacts of Roads: The Case for Road Density as Indicator of Human Disturbance and Road-Density Reduction as Restoration Target; A Concise Review. Pacific Rivers Council Science Publication 09-001. Pacific Rivers Council, Portland, OR and Polson, MT. Available at: <http://www.pacificrivers.org/science-research/resources-publications/road-density-as-indicator/download>
- Coffin, A. 2006. From roadkill to road ecology: A review of the ecological effects of roads. *Journal of Transport Geography* 15: 396-406.
- Crist, M.R., and B. Wilmer. 2002. *Roadless Areas: The Missing Link in Conservation*. The Wilderness Society, Washington D.C.

- Davenport, J., and T.A. Switalski. 2006. Environmental impacts of transport related to tourism and leisure activities. In: *The ecology of transportation: managing mobility for the environment*, editors: J Davenport and Julia Davenport. Dordrecht, Netherlands: Kluwer Academic Publishers. 333-360. Available at: [http://www.wildlandscpr.org/files/uploads/PDFs/d\\_Switalski\\_2006\\_Enviro\\_impacts\\_of\\_transport.pdf](http://www.wildlandscpr.org/files/uploads/PDFs/d_Switalski_2006_Enviro_impacts_of_transport.pdf)
- DellaSala, D., J. Karr, and D. Olson. 2011. Roadless areas and clean water. *Journal of Soil and Water Conservation*, vol. 66, no. 3. May/June 2011.
- DeVelice, R., and J.R. Martin. 2001. Assessing the extent to which roadless areas complement the conservation of biological diversity. *Ecological Applications* 11(4): 1008-1018.
- Endicott, D. 2008. National Level Assessment of Water Quality Impairments Related to Forest Roads and Their Prevention by Best Management Practices. A Report Prepared by the Great Lakes Environmental Center for the Environmental Protection Agency, Office of Water, December 4, 2008. 259 pp.
- Edge, W.D., and C.L. Marcum. 1985. Movements of elk in relation to logging disturbances. *Journal of Wildlife Management* 49(4): 926–930.
- Fahrig, L., and T. Rytwinski. 2009. Effects of roads on animal abundance: an empirical review and synthesis. *Ecology and Society* 14(1): 21.  
Available at: <http://www.ecologyandsociety.org/vol14/iss1/art21/>.
- Foltz, R.B. N.S. Copeland, and W.J. Elliot. 2009. Reopening abandoned forest roads in northern Idaho, USA: Quantification of runoff, sediment concentration, infiltration, and interrill erosion parameters. *Journal of Environmental Management* 90: 2542–2550.
- Forman, R. T. T., and A.M. Hersperger. 1996. Road ecology and road density in different landscapes, with international planning and mitigation solutions. Pages 1–22. IN: G. L. Evinck, P. Garrett, D. Zeigler, and J. Berry (eds.), *Trends in Addressing Transportation Related Wildlife Mortality*. No. FLER- 58-96, Florida Department of Transportation, Tallahassee, Florida.
- Foreman, R.T.T., D. Sperling, J.A. Bissonette et al. 2003. *Road Ecology – Science and Solutions*. Island Press. Washington, D.C. 504 p.
- Furniss, M.J., T.D. Roelofs, and C.S. Yee. 1991. Road construction and maintenance. In: Meehan, W.R., ed. *Influences of forest and rangeland management on salmonid fishes and their habitats*. Spec. Publ. 19. Bethesda, MD: American Fisheries Society. p. 297-323.
- Gaines, W.L., P. Singleton, and R.C. Ross. 2003. Assessing the cumulative effects of linear recreation routes on wildlife habitats on the Okanogan and Wenatchee National Forests. Gen. Tech. Rep. PNW-GTR-586. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 79 p. Available at: <http://www.montanawildlife.com/projectsissues/Assessingthecumulativeeffectsoflinearrecreationroutesonwildlifehabitats.pdf>

- Gelbard, J.L., and S. Harrison. 2003. Roadless habitats as refuges for native grasslands: interactions with soil, aspect, and grazing. *Ecological Applications* 13(2): 404-415.
- Girvetz, E., and F. Shilling. 2003. Decision Support for Road System Analysis and Modification on the Tahoe National Forest. *Environmental Management* 32(2): 218–233
- Grant, A., C.R. Nelson, T.A. Switalski, and S.M. Rinehart. 2011. Restoration of native plant communities after road decommissioning in the Rocky Mountains: effect of seed mix composition & soil properties on vegetative establishment. *Restoration Ecology* 19: 160-169.
- Gucinski, M., J. Furniss, R. Ziemer, and M.H. Brookes. 2000. *Forest Roads: A Synthesis of Scientific Information*. Gen. Tech. Rep. PNWGTR-509. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 103 p. Available at: <http://www.fs.fed.us/pnw/pubs/gtr509.pdf>.
- Hargis, C.D., J.A. Bissonette, and D.T. Turner. 1999. The influence of forest fragmentation and landscape pattern on American martens. *Journal of Applied Ecology* 36(1): 157–172.
- Hebblewhite, M., R.H. Munro, E.H. Merrill. 2009. Trophic consequences of postfire logging in a wolf-ungulate system. *Forest Ecology and Management* 257(3): 1053-1062.
- Holman, I.P., R.J. Nicholls, P.M. Berry, P.A. Harrison, E. Audsley, S. Shackley, and M.D.A. Rounsevell. 2005. A regional, multi-sectoral and integrated assessment of the impacts of climate and socio-economic change in the UK. Part II. Results. *Climatic Change* 71: 43-73.
- Jalkotzy, M.G., P.I. Ross, and M.D. Nasserden. 1997. The effects of linear developments on wildlife: a review of selected scientific literature. Prepared for Canadian Association of Petroleum Producers. Arc Wildlife Services, Ltd., Calgary, AB. 115 p.
- Jensen W.F., T.K. Fuller, and W.L. Robinson. 1986. Wolf (*Canis lupus*) distribution on the Ontario-Michigan border near Sault Ste. Marie. *Canadian Field-Naturalist* 100: 363-366.
- Joslin, G., and H. Youmans, coordinators. 1999. Effects of recreation on Rocky Mountain wildlife: A Review for Montana. Committee on Effects of Recreation on Wildlife, Montana Chapter of The Wildlife Society. 307 p. Available at: <http://joomla.wildlife.org/Montana/index>
- Kasworm, W.F., and T.L. Manley. 1990. Road and trail influences on grizzly bears and black bears in northwest Montana. *International Conference on Bear Research and Management* 8: 79-84.
- Kerkvliet, J., J. Hicks, and B. Wilmer. 2008. Carbon Sequestered when Unneeded National Forest Roads are Revegetated. The Wilderness Society Briefing Memo. Available at: [http://wilderness.org/sites/default/files/legacy/brief\\_carbonandroads.pdf](http://wilderness.org/sites/default/files/legacy/brief_carbonandroads.pdf).
- Lee, D., J. Sedell, B.E. Rieman, R. Thurow, and J. Williams. 1997. Broad-scale assessment of aquatic species and habitats. In: An assessment of ecosystem components in the interior

Columbia basin and portions of the Klamath and Great Basins. Edited by T.M. Quigley and S.J. Arbelvide. General Technical Report PNW-GTR-405. USDA Forest Service, Pacific Northwest Research Station, Portland, OR. Vol III. p. 183–196.

Lloyd, R., K. Lohse, and T.P.A. Ferre. 2013. Influence of road reclamation techniques on forest ecosystem recovery. *Frontiers in Ecology and the Environment* 11(2): 75-81.

Loucks, C., N. Brown, A. Loucks, and K. 2003. USDA Forest Service roadless areas: potential biodiversity conservation reserves. *Conservation Ecology* 7(2): 5.  
Available at: <http://www.ecologyandsociety.org/vol7/iss2/art5/>

Logan, R. 2001. Water Quality BMPs for Montana Forests. Montana Department of Environmental Quality. Missoula, MT. 60p. Available at:  
<https://dnrc.mt.gov/Forestry/Assistance/Practices/Documents/2001WaterQualityBMPGuide.pdf>

Lyon, L.J. 1983. Road density models describing habitat effectiveness for elk. *Journal of Forestry* 81: 592-595.

Mace, R.D., J.S. Waller, T.L. Manley, L.J. Lyon, and H. Zuuring. 1996. Relationships among grizzly bears, roads and habitat in the Swan Mountains, MT. *Journal of Applied Ecology*. 33: 1395-1404.

Madej, M., J. Seney, and P. van Mantgem. 2013. Effects of road decommissioning on carbon stocks, losses, and emissions in north coastal California. *Restoration Ecology* 21(4): 439–446.

Mattson, D.J., S. Herrero, R.G. Wright, and C.M. Pease. 1996. Science and management of Rocky Mountain grizzly bears. *Conservation Biology* 10(4): 1013-1025.

McCaffery M., T.A. Switalski, and L. Eby. 2007. Effects of road decommissioning on stream habitat characteristics in the South Fork Flathead River, Montana. *Transactions of the American Fisheries Society* 136: 553-561.

McGurk, B.J., and D.R. Fong, 1995. Equivalent roaded area as a measure of cumulative effect of logging. *Environmental Management* 19: 609-621.

Mech, L D. 1989. Wolf population survival in an area of high road density. *American Midland Naturalist* 121: 387-389.

Mech, L. D., S.H. Fritts, G.L. Radde, and W.J. Paul. 1988. Wolf distribution and road density in Minnesota. *Wildlife Society Bulletin* 16: 85-87.

Merrill, B.R., and E. Cassaday. 2003. Best Management Practices for Road Rehabilitation – Road – Stream Crossing Manual. California State Parks. Eureka, CA. 25p. Available at:  
[http://www.parks.ca.gov/pages/23071/files/streamcrossingremovalbmp5\\_03.pdf](http://www.parks.ca.gov/pages/23071/files/streamcrossingremovalbmp5_03.pdf)

- Mladenoff, D.J., T.A. Sickley, R.G. Haight, and A.P. Wydeven. 1995. A regional landscape analysis and prediction of favorable gray wolf habitat in the Northern Great Lakes region. *Conservation Biology* 9: 279-294.
- Moore, T. 2007. [unpublished draft]. National Forest System Road Trends, Trends Analysis Submitted to Office of Management and Budget. United States Department of Agriculture, Forest Service, Engineering Staff, Washington Office, Washington, DC.
- National Fish, Wildlife and Plants Climate Adaptation Partnership (NFWPCAP). 2012. National Fish, Wildlife and Plants Climate Adaptation Strategy. Association of Fish and Wildlife Agencies, Council on environmental Quality, Great Lakes Indian Fish and Wildlife Commission, National Oceanic and Atmospheric Administration, and U.S. Fish and Wildlife Service. Washington, DC.
- Noss, R.F. 2001. Beyond Kyoto: forest management in a time of rapid climate change. *Conservation Biology* 15(3): 578-590.
- Ortega, Y.K., and D.E. Capen. 2002. Roads as edges: effects on birds in forested landscapes. *Forest Science* 48(2): 381-396.
- Ouren, D.S., C. Haas, C.P. Melcher, S.C. Stewart, P.D. Ponds, N.R. Sexton, L. Burris, T. Fancher, and Z.H. Bowen. 2007. Environmental effects of off-highway vehicles on Bureau of Land Management lands: A literature synthesis, annotated bibliographies, extensive bibliographies, and internet resources: U.S. Geological Survey, Open-File Report 2007-1353, 225 p.
- Parmesan, C. 2006. Ecological and evolutionary responses to recent climate change. *Annual Review of Ecology, Evolution, and Systematics* 37: 637-669.
- Quigley, T.M., and S.J. Arbelbide, tech. eds. 1997. An assessment of ecosystem components in the interior Columbia basin and portions of the Klamath and Great Basins: volume 1 and volume 3. Gen. Tech. Rep. PNW-GTR-405. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station.  
Available at: [http://www.fs.fed.us/pnw/publications/pnw\\_gtr405/](http://www.fs.fed.us/pnw/publications/pnw_gtr405/).
- Reynolds, K. 1999. Netweaver for EMDS user guide (version1.1); a knowledge base development system. General technical Report PNW-GTR-471. USDA Forest Service, Pacific Northwest Research Station, Portland, OR.
- Rhodes, J.J., McCullough, D.A., and F.A. Espinosa. 1994. A coarse screening process for evaluation of the effects of land management activities on salmon spawning and rearing habitat in ESA consultations. Tech. Rep. 94-4. Portland, OR: Columbia River Intertribal Fish Commission. 127 p.
- Rieman, B., D. Lee, G. Chandler, and D. Myers. 1997. Does wildfire threaten extinction for salmonids? Responses of Redband Trout and Bull Trout Following Recent Large Fires on the Boise National Forest, in Greenlee, J. M., Proceedings: First Conference on Fire Effects on



Rare and Endangered Species and Habitats. Coeur d'Alene, Idaho. International Association of Wildland Fire. Fairfield, WA. p. 47-57.

- Robichaud, P.R., L.H. MacDonald, and R.B. Foltz. 2010. Fuel management and Erosion. In: Cumulative Watershed Effects of Fuels Management in the Western United States. USDA Forest Service RMRS-GTR-231. P. 79-100. Available at: [http://www.fs.fed.us/rm/pubs/rmrs\\_gtr231/rmrs\\_gtr231\\_079\\_100.pdf](http://www.fs.fed.us/rm/pubs/rmrs_gtr231/rmrs_gtr231_079_100.pdf)
- Robinson, C., P.N. Duinker, and K.F. Beazley. 2010. A conceptual framework for understanding, assessing, and mitigation effects for forest roads. *Environmental Review* 18: 61-86.
- Rooney, T.P. 2006. Distribution of ecologically-invasive plants along off-road vehicle trails in the Chequamegon National Forest, Wisconsin. *The Michigan Botanist* 44:178-182
- Rost, G.R., and J.A. Bailey. 1979. Distribution of mule deer and elk in relation to roads. *Journal of Wildlife Management* 43(3): 634-641.
- Rowland, M.M., M.J. Wisdom, B.K. Johnson, and M.A. Penninger. 2005. Effects of roads on elk: implications for management in forested ecosystems. Pages 42-52. IN: Wisdom, M.J., technical editor, *The Starkey Project: a Synthesis of Long-term Studies of Elk and Mule Deer*. Reprinted from the 2004 Transactions of the North American Wildlife and Natural Resources Conference, Alliance Communications Group, Lawrence, KS.
- Schiffman, L. 2005. Archaeology, Off-Road Vehicles, and the BLM. Published online April 20, 2005. *Archeaology*. Available at: <http://www.archaeology.org/online/features/southwest/>
- Semlitsch, R.D., T.J. Ryan, K. Hamed, M. Chatfield, B. Brehman, N. Pekarek, M. Spath, and A. Watland. 2007. Salamander abundance along road edges and within abandoned logging roads in Appalachian forests. *Conservation Biology* 21: 159-167.
- Shilling, F., J. Boggs, and S. Reed. 2012. Recreational System Optimization to Reduce Conflict on Public Lands. *Environmental Management* 50: 381-395.
- Strittholt, J., and D. Dellasala. 2001. Importance of Roadless Area Conservation in Forested Ecosystems: Case Study of the Klamath-Siskiyou Region of the United States. In *Conservation Biology* 15(6): 1742-1754.
- Switalski, T.A., J.A. Bissonette, T.H. DeLuca, C.H. Luce, and M.A. Madej. 2004. Benefits and impacts of road removal. *Frontiers in Ecology and the Environment*. 2(1): 21-28. Available at: [http://www.fs.fed.us/rm/pubs\\_other/rmrs\\_2004\\_switalski\\_t001.pdf](http://www.fs.fed.us/rm/pubs_other/rmrs_2004_switalski_t001.pdf)
- Switalski, T.A., and C.R. Nelson. 2011. Efficacy of road removal for restoring wildlife habitat: black bear in the Northern Rocky Mountains, USA. *Biological Conservation* 144: 2666-2673.

- Switalski, T.A., and A. Jones. 2012. Off-road vehicle best management practices for forestlands: A review of scientific literature and guidance for managers. *Journal of Conservation Planning* 8: 12-24.
- The Wilderness Society. 2004. *Landscape Connectivity: An Essential Element of Land Management*. Policy Brief. Number 1.
- The Wilderness Society. 2012. *Rightsizing the National Forest Road System: A Decision Support Tool*. Available at: <http://www.landscapecollaborative.org/download/attachments/12747016/Road+decommissioning+model+overview+2012-02-29.pdf?version=1&modificationDate=1331595972330>.
- The Wilderness Society. 2013. *RoadRight: A Spatial Decision Support System to Prioritize Decommissioning and Repairing Roads in National Forests User Guide*. RoadRight version: 2.2, User Guide version: February, 2013. Available at: <http://www.landscapecollaborative.org/download/attachments/18415665/RoadRight%20User%20Guide%20v22.pdf?api=v2>
- Thiel, R.P. 1985. The relationships between road densities and wolf habitat in Wisconsin. *American Midland Naturalist* 113: 404-407.
- Trombulak S., and C. Frissell. 2000. Review of Ecological Effects of Roads on Terrestrial and Aquatic Communities. *Conservation Biology* 14(1): 18-30.
- USDA Forest Service. 1996a. *National Forest Fire Report, 1994*. Washington DC.
- USDA Forest Service. 1996b. *Status of the interior Columbia basin: summary of scientific findings*. Gen. Tech. Rep. PNW-GTR-385. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station; U.S. Department of the Interior, Bureau of Land Management. 144 p.
- USDA Forest Service. 1998. *1991-1997 Wildland Fire Statistics*. Fire and Aviation Management, Washington, D.C.
- USDA Forest Service. 1999. *Roads Analysis: Informing Decisions about Managing the National Forest Transportation System*. Misc. Rep. FS-643. Washington, D.C.: USDA Forest Service. 222 p. Available at: [http://www.fs.fed.us/eng/road\\_mgt/DOCSroad-analysis.shtml](http://www.fs.fed.us/eng/road_mgt/DOCSroad-analysis.shtml)
- USDA Forest Service. 2001a. *Final National Forest System Road Management Strategy Environmental Assessment and Civil Rights Impact Analysis*. U.S. Department of Agriculture Forest Service Washington Office, January 2001.
- USDA Forest Service. 2010. *Water, Climate Change, and Forests: Watershed Stewardship for a Changing Climate*, PNW-GTR-812, June 2010, 72 p. Available at: [http://www.fs.fed.us/pnw/pubs/pnw\\_gtr812.pdf](http://www.fs.fed.us/pnw/pubs/pnw_gtr812.pdf).

- USDA Forest Service. 2011a. Adapting to Climate Change at Olympic National Forest and Olympic National Park. Forest Service Pacific Northwest Research Station General Technical Report, PNW-GTR-844, August 2011.  
Available at: [http://www.fs.fed.us/pnw/pubs/pnw\\_gtr844.pdf](http://www.fs.fed.us/pnw/pubs/pnw_gtr844.pdf)
- USDA Forest Service. 2011b. National Roadmap for Responding to Climate Change. US Department of Agriculture. FS-957b. 26 p.  
Available at: [http://www.fs.fed.us/climatechange/pdf/Roadmap\\_pub.pdf](http://www.fs.fed.us/climatechange/pdf/Roadmap_pub.pdf).
- USDA Forest Service. 2012a. Assessing the Vulnerability of Watersheds to Climate Change: Results of National Forest Watershed Vulnerability Pilot Assessments. Climate Change Resource Center.
- USDA Forest Service. 2012b. National Best Management Practices for Water Quality Management on National Forest System Lands. Report# FS-990. 177p. Available at: [http://www.fs.fed.us/biology/resources/pubs/watershed/FS\\_National\\_Core\\_BMPs\\_April2012.pdf](http://www.fs.fed.us/biology/resources/pubs/watershed/FS_National_Core_BMPs_April2012.pdf)
- USDI Fish and Wildlife Service. 1993. Grizzly bear recovery plan. Missoula, MT. 181p.
- USDI Fish and Wildlife Service. 1999. Endangered and Threatened Wildlife and Plants; Determination of Threatened Status for Bull Trout in the Coterminous United States; Final Rule. Federal Register Volume 64, Number 210 (Monday, November 1, 1999). p. 58922.
- USDI Bureau of Land Management. 2000. Strategic paper on cultural resources at risk. Bureau of Land Management, Washington, D.C. 18 p.
- USDI National Park Service. 2010. Climate Change Response Strategy. National Park Service Climate Change Response Program, Fort Collins, Colorado.  
Available at: [http://www.nature.nps.gov/climatechange/docs/NPS\\_CCRS.pdf](http://www.nature.nps.gov/climatechange/docs/NPS_CCRS.pdf).
- van Dyke, F.G., R.H. Brocke, H.G. Shaw, B.B Ackerman, T.P. Hemker, and F.G. Lindzey. 1986. Reactions of mountain lions to logging and human activity. *Journal of Wildlife Management*. 50(1): 95–102.
- Wasser, S.K., K. Bevis, G. King, and E. Hanson. 1997. Noninvasive physiological measures of disturbance in the northern spotted owl. *Conservation Biology* 11(4): 1019–1022.
- Wemple, B.C., J.A. Jones, and G.E. Grant. 1996. Channel network extension by logging roads in two basins, western Cascades, OR. *Water Resources Bulletin* 32: 1195-1207.
- Wemple, B.C., F.J. Swanson, and J.A. Jones. 2001. Forest Roads and geomorphic process interactions, Cascade Range, Oregon. *Earth Surface Process and Landforms* 26: 191-204.  
Available at: <http://andrewsforest.oregonstate.edu/pubs/pdf/pub2731.pdf>
- Wilcove, D.S. 1990. The role of wilderness in protecting biodiversity. *Natural Resources and Environmental Issues*: Vol. 0, Article 7.

Wisdom, M.J., R.S. Holthausen, B.C. Wales, C.D. Hargis, V.A. Saab, D.C. Lee, W.J. Hann, T.D. Rich, M.M. Rowland, W.J. Murphy, and M.R. Eames. 2000. Source habitats for terrestrial vertebrates of focus in the interior Columbia basin: Broad-scale trends and management implications. Volume 1 – Overview. Gen. Tech. Rep. PNW-GTR-485. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station.

Wydeven, A.P, D.J. Mladenoff, T.A. Sickley, B.E. Kohn, R.P. Thiel, and J.L. Hansen. 2001. Road density as a factor in habitat selection by wolves and other carnivores in the Great Lakes Region. *Endangered Species Update* 18(4): 110-114.

### Attachments

Attachment 1: Wildfire and Roads Fact Sheet

Attachment 2: Using Road Density as a Metric for Ecological Health in National Forests: What Roads and Routes should be Included? Summary of Scientific Information